

Table 1) Frequency of demographic characteristics in patients with spinal cord injury (n=248)

Parameter	Number	Percentage
Gender		
Female	27	10.89
Male	221	89.11
Age Range		
Less than 30	32	12.9
30-40 Years	106	42.74
40-50Years	85	34.27
50-60 Years	25	10.08
Marital Status		
Single	57	22.99
Married	191	77.01
Education Level		
Less than Diploma	87	35.08
Diploma	103	41.53
University Degree	58	23.39
Cause		
War	184	74.19
Accident with Vehicle	48	19.35
Spinal Cord Tumors	4	1.61
Fall from Height	12	4.83
The Type of Injury		
Quadriplegia	16	6.46
Paraplegia	232	93.54

Table 2) Prevalence and intensity of pain in patients with spinal cord injury

Parameter	Prevalence	Intensity of Pain	Range
Without Pain	6 (2.4)	-	-
Neuropathic Pain	204 (82.3)	8.12±1.72	6-10
Musculoskeletal Pain	201 (81.0)	6.06±1.22	4-9
Visceral Pain	102 (41.1)	3.88±0.89	2-7
Other Pains	3 (1.2)	5.66±0.57	5-6

Table 3) Frequency of involvement pain location in patients with spinal cord injury

Location of Pain	Neuropathic Pain	Musculoskeletal Pain	Visceral Pain
Head	4	-	-
Neck and Shoulder	38	168	-
Arm and Hand	25	172	-
Torso and Genitals	16	27	88
Back	46	78	39
Buttocks and Hips	40	-	-
Thigh	48	-	-
Leg and Foot	72	-	-