

**Table 1)** Relative and absolute frequency (numbers in parentheses are percentage) of IUD types used in women referring to Sarem Medical Center between 1985 and 1995 (n=1199)

IUD Type	Frequency
Copper7	200(16.68)
Copper T	599(49.96)
T380A	200(16.68)
Multiload	200(16.68)

**Table 2)** Absolute and relative frequency (number in parenthesis are percentage) of IUD displacement cases, one month after placement in women referred to Sarem Medical Center between 1985 and 1995 (n=1199)

Status of IUD Placement	Frequency
Normal	1179(98.30)
Displaced	20(1.70)

**Table 3)** Frequency rate of failure of various contraceptive methods after 5 years in different countries

Country	Tablet	IUD	Periodical	Natural	Other	All Methods Include Sterilization	All Methods without Sterilization
<b>Latin America</b>							
Bolivia	5.10	0.30	2.21	8.34	4.35	2.19	4.20
Brazil	4.50	0.13	8.22	7.20	8.16	9.70	4.10
Columbia	7.70	3.50	8.24	1.20	6.23	7.11	7.13
Dominican	8.11	6.30	9.19	2.29	3.23	7.80	0.14
Ecuador	8.50	6.40	9.13	2.27	1.21	1.80	0.10
Guatemala	8.90	7.70	6.24	0.12	6.16	9.10	5.14
Mexico	4.50	5.10	7.30	0.40	8.50	2.30	1.40
Peru	9.50	2.40	6.23	3.17	8.16	7.15	7.16
Trinidad and Tobago	9.50	2.40	7.13	6.17	2.15	3.10	3.11
<b>Asia</b>							
China	5.14	3.10	8.15	9.15	7.15	2.70	3.11
Indonesia	7.20	9.10	8.10	5.80	6.20	8.20	9.20
Sri Lanka	2.70	4.30	0.13	5.11	1.80	4.70	6.10
Thailand	8.20	5.20	8.14	4.70	1.10	9.10	5.20
<b>North Africa</b>							
Egypt	9.60	8.10	4.38	3.10	9.8	0.6	0.60
Morocco	6.80	1.10	5.30	5.11	6.22	9.10	1.11
Tunisia	4.50	7.20	2.14	5.12	2.21	4.70	0.80