

Table 1) The Frequency of demographic characteristics of veterans with neuro-musculoskeletal disorders

Parameter	Frequency	Percentage
Gender		
Male	786	98.9
Female	9	1.1
Age Group		
30 Years and less	5	0.6
31-40 Years	21	2.6
41-50 Years	345	43.4
51-60 Years	325	40.9
61-70 Years	74	9.3
71 Years and more	25	3.1
Percentage of injury		
25% and less	358	45.0
26-49%	338	42.5
50% and more	99	12.5
Marital status		
Never being married	7	0.9
Married	779	98.0
Divorced or widower	9	1.1
Number of Children		
3 Children and less	492	61.9
4 Children and more	303	38.1
Veterans' number of family members		
3 and less	41	5.2
4-6	228	28.7
7-9	350	44.0
10 and more	176	22.1
Employment Status		
Employed	309	38.9
Unemployed	486	61.1
Education at the time of study		
Illiterate	51	6.4
Below Diploma	274	34.5
Diploma	234	29.4
Higher than diploma	236	29.7
Habitat Province		
Zanjan	81	10.2
Markazi	136	17.1
Isfahan	114	14.3
Hamedan	140	17.6
West Azerbaijan	142	17.9
Mazandaran	40	5.0
Ardebil	24	3.0
Kermanshah	118	14.8

Table 2) Frequency of the causes of injuries and types of accompanying injury in the veterans with leg and ankle neuro-musculoskeletal disorders

Parameter	Frequency	Percentage
Cause of Injury		
Quiver	397	49.9
Shot	181	22.8
Landmine	119	15.0
Blast wave	63	7.9
Others	35	4.4
Accompanying injury*		
Neurology and psychiatry	253	18.2
Chemical	91	6.6
Spinal cord injury	28	2.0
Head injury	153	11.0
Face injury	107	7.7
Chest Injury	76	5.5
Back and spine injury	161	11.6
Abdominal injury	135	9.7
Upper limb injury	221	15.9
Others	164	11.8

*Some veterans had more than one accompanying injury. Therefore, the total number of accompanying injuries is more than sample size

Table 3) Absolute and relative frequency (numbers in the parentheses are based on percentage) of different injuries in the veterans with ankle and leg neuro-musculoskeletal

Type of Injury	Right	Left	Without Disorders
Limitation of ankle dorsiflexion range of motion	282 (35.5)	259(32.6)	254(31.9)
Limb shortening	302(38.0)	285(35.8)	208(26.7)
Limb amputation	17(2.2)	13(1.6)	765(96.2)
Lower Limbs Peripheral nerve injuries			
Sciatic	81(10.7)	61(7.7)	653(82.1)
Femoral	15(1.9)	15(1.9)	765(96.2)
Peroneal	79(9.9)	83	633
Tibial	64(8.1)	60(7.5)	671(84.4)