

Table 1) Absolute and relative frequency (numbers in parentheses are percentage) of anatomic location of amputated organ in amputee veterans participated in the study (n=256)

Location of Amputation	Right	Left	Total
Upper Limbs			
Shoulder	18(7.2)	17(6.6)	35(13.8)
Above the Elbow	15(5.9)	10(3.9)	25(9.8)
On the Elbow	6(2.3)	3(1.2)	9(3.5)
Below the Elbow	8(3.1)	8(3.1)	16(2.6)
Wrist	7(2.7)	4(1.6)	11(4.3)
Hand	5(2.0)	2(0.8)	7(2.8)
Fingers	7(2.7)	8(3.1)	15(5.8)
Total	74(25.9)	52(20.3)	126(46.2)
Lower Limbs			
Hemicolectomy	0	5(2.0)	5(2.0)
Hip	11(4.3)	7(2.7)	17(0.7)
The Upper Third of Thigh, above the knee	18(7.0)	14(5.5)	32(12.5)
Mid-thigh, above the knee	9(3.5)	6(2.3)	15(5.8)
The Lower Third of the Thigh, above the Knee	9(3.5)	10(3.9)	19(7.4)
Knee	8(3.1)	7(2.7)	15(5.8)
Very close to the Knee, Below the Knee	28(10.9)	19(7.4)	47(18.3)
A third Upper leg, Below the Knee	18(7.0)	11(4.3)	29(11.3)
Middle of the Upper Third of the Leg, Below the Knee	12(4.7)	16(6.2)	28(10.9)
The Lower Third of the Leg below the Knee	4(1.6)	6(2.3)	10(3.9)
Ankle	23(9.0)	23(9.0)	46(18.0)
Toe	3(1.2)	4(1.6)	7(2.8)
Toe Finger	1(0.4)	2(0.8)	3(1.2)
Total	144(56.2)	130(50.7)	274(106.9)
Upper and Lower Limbs			
Right Upper and Lower Limbs	-----	-----	10(3.9)
Left Upper and Lower Limbs	-----	-----	2(0.8)
Right and Left Upper Limb and Right Lower Limb	-----	-----	4(1.6)
Right and Left Upper Limb and Left Lower Limb	-----	-----	6(2.3)
Right and Left Lower Limb and Right Upper Limb	-----	-----	8(3.1)
Right and Left Lower Limb with Left Upper Limb	-----	-----	6(2.3)
Left Upper Limb and Right Lower Limb	-----	-----	5(2.0)
Right Upper Limb and Left Lower Limb	-----	-----	3(1.20)

Table 2) The frequency of clinical complications in veterans with amputations participating in the study (n=256)

Clinical Complications	Number	Percentage
Cook Syndrome	5	2.0
Skin Infection	25	9.8
Skin Inflammation	29	11.3
Bone Spur	18	7.0
Pain	105	41.0
Phantom Pain	43	16.8
Trauma	11	4.3
Phantom Feeling	34	13.3
Contraction	41	16.0

Table 3) Mean score of nine scales and three indices of SCL-90-R in amputee veterans who participated in the study

Total Subjects (n=256)	Probable Patient (n=116)	Probable Healthy Person (n=140)
Physical Complaints		
1.60±0.86	2.27±0.62	1.05±0.61
Obsessive Compulsive Disorders		
1.33±0.86	2.10±0.59	0.7±0.44
Sensitivity in mutual Relations		
1.17±0.89	1.93±0.68	0.54±0.43
Depression		
1.29±0.93	2.11±0.68	0.61±0.41
Anxiety		
1.34±0.89	2.11±0.64	0.7±0.47
Aggression		
1.25±0.91	2.00±0.71	0.63±0.51
Phobia		
0.86±0.81	1.40±0.75	0.35±0.34
Paranoid Thoughts		
1.28±0.90	1.92±0.80	0.70±0.58
Psychosis		
0.84±0.73	1.41±0.66	0.35±0.34
Global Severity Index (GSI)		
1.24±0.76	1.94±0.49	0.66±0.34
Positive Symptom Total (PST)		
50.98±23.10	71.10±11.00	34.20±15.00
Positive Symptom Distress Index (PSDI)		
2.03±0.60	2.46±0.40	1.67±0.46