

Table 1) Distribution of absolute frequency of demographic characteristics in the group of people with drug abuse disorder and normal people (each group of 100 persons)

Demographic Variables	People with Drug Abuse Disorder	Normal People
Marital Status		
Married	53	41
Single	47	59
Level of Education		
Junior High School	29	25
Diploma	36	32
Associate	16	20
Bachelor	13	15
Master	6	8

Table 2) Distribution of absolute frequency of type of drug and daily intake in people with drug abuse disorder (100 persons)

Variables	Number
Type of Drug	
Opium	17
Cannabis	3
Heroin	27
Methadone	6
Tablet	5
Glass	22
Cocaine	2
Alcohol	1
Multi-Type	17
Daily Intake	
Less than a gram	42
1-2 gram	40
2 gram and over	18

Table 3) Comparison of the mean scores of hypersexual disorder and its dimensions in people with drug abuse disorder (100 persons) and normal people (100 people)

Variables	People with Drug Abuse Disorder	Normal People	F Value	Differentiated Eta Coefficient
Hypersexual Disorder				
Mean	81.86±14.89	42.80±11.96	30.41	0.13
Range	21-90	19-67		
Copying Dimension				
Mean	19.86±7.16	14.29±5.73	36.82	0.15
Range	7-32	7-31		
Consequences Dimension				
Mean	11.18±4.55	8.55±2.95	23.47	0.10
Range	4-20	4-17		
Control Dimension				
Mean	23.85±8.32	19.96±6.02	14.32	0.06
Range	8-40	8-32		

In all cases $p < 0.001$