

Table 1. The mean and range of measured variables in patients with oral opium consume referring with abdominal pain (n=122)

Variables	Mean	Min	Max
Blood lead level ($\mu\text{g}/\text{dl}$)	87.49 \pm 32.27	0.26	182.0
Hemoglobin (g / dl)	10.94 \pm 1.96	0.7	18.70
Hematocrit (%)	33.19 \pm 5.41	50.22	54.20
MCV (femtoliter)	81.91 \pm 8.98	70.33	96.60
SGPT (IU/L)	49.76 \pm 15.23	00.5	324.00

Table 2. The absolute and relative frequency of research variables at different levels in patients with oral opium consume referring with abdominal pain (n = 122)

Variables	No.	%
Blood lead level (µg/dl)		
Below 25 (non-toxic)	0	0
25-45 (toxic)	4	3.3
45-69 (Mild toxicity)	36	29.5
99-70 (moderate toxicity)	48	39.3
More than 100 (severe toxicity)	34	27.9
Hemoglobin (g / dl)		
More than 12	29	23.8
10-12	52	42.6
8-9.9	34	27.9
Below 8	7	5.7
Hematocrit (%)		
Below 40	100	82.0
40-54	20	16.4
More than 54	2	1.6
MCV (femtoliter)		
Below 80	32	26.2
96-80	88	72.1
More than 96	2	1.6
SGPT (IU/L)		
Below 45	65	53.3
45-89	33	27.0
90-134	18	14.8
135-180	0	0
More than 189	6	4.9